

Not to be cited without
permission of the authors¹

Canadian Atlantic Fisheries
Scientific Advisory Committee

CAFSAC Research Document 87/97

Ne pas citer sans
autorisation des auteurs¹

Comité scientifique consultatif des
pêches canadiennes dans l'Atlantique

CSCPCA Document de recherche 87/97

**An evaluation of the population dynamics of 4X haddock
during 1962-87 with yield projected to 1988.**

by

R. O'Boyle and D. Wallace
Marine Fish Division
Fisheries and Oceans
P.O. Box 1006
Dartmouth, N.S. B2Y 4A2

¹ This series documents the scientific basis for fisheries management advice in Atlantic Canada. As such, it addresses the issues of the day in the time frames required and the Research Documents it contains are not intended as definitive statements on the subjects addressed but rather as progress reports on ongoing investigations.

Research Documents are produced in the official language in which they are provided to the Secretariat by the author.

¹ Cette série documente les bases scientifiques des conseils de gestion des pêches sur la côte atlantique du Canada. Comme telle, elle couvre les problèmes actuels selon les échéanciers voulus et les Documents de recherche qu'elle contient ne doivent pas être considérés comme des énoncés finals sur les sujets traités mais plutôt comme des rapports d'étape sur les études en cours.

Les Documents de recherche sont publiés dans la langue officielle utilisée par les auteurs dans le manuscrit envoyé au secrétariat.

Abstract

This evaluation of the 4X haddock population is similar to that conducted last year. The calibration models are virtually identical. Only the input data have changed. In this regard the two assessments are very different, with the 1987 survey data providing a pessimistic view of resource status. The 1986 fully recruited fishing mortality is estimated to be 0.513 with a comparable level (0.549) generated by the 15,000 t TAC in 1987. Total stock biomass is currently about 60,000 t, reaching the low levels observed in the early 1970's. Poor recruiting 1984 - 86 year-classes are expected to push this biomass and consequent yield even lower over the next 3 to 5 years. The resource is currently suffering from growth overfishing and recruitment overfishing is highly likely. Resource projections to 1988 indicate an $F_{0.1}$ yield of 8204 t while the 50 percent rule level is 12,384 t. Given market pressure, these levels will likely be exceeded.

Résumé

La présente évaluation de la population d'aiglefin 4X est similaire à celle effectuée l'an dernier. Les modèles d'étalonnage sont pratiquement identiques. Seules les données d'entrée ont changé. A cet égard, les deux évaluations sont très différentes, les données du relevé de 1987 donnant une vue pessimiste de l'état de la ressource. On estime à 0,513 la mortalité par pêche du poisson entièrement recruté en 1986 et le TPA de 15 000 t pour 1987 donne un niveau comparable (0,549). Actuellement, la biomasse totale du stock est d'environ 60 000 t, retombant ainsi aux bas niveaux observés au début des années 70. On prévoit que le faible recrutement des classes de 1984 à 1986 va encore faire baisser la biomasse, et partant, le rendement, au cours des 3 à 5 prochaines années. La ressource souffre actuellement d'une surpêche au détriment de la croissance et il est très probable que la surpêche nuise au recrutement. Les prévisions pour la ressource jusque 1988 donnent un rendement à $F_{0.1}$ de 8 204 t tandis que le niveau avec la règle de 50 pour cent est de 12 384 t. Compte tenu de la pression du marché, ces niveaux seront probablement dépassés.

Introduction

Sequential Population Analysis (SPA) has been used to provide estimates of the size of the haddock population in NAFO Division 4X since 1974. These analyses have been calibrated using groundfish survey data. Since 1980, US survey sampling in NAFO Division 4X was restricted to those strata covering Brown's Bank. Commercial catch rate indices have also been used, although their utility has been limited by problems in data quality. Consequently, in recent years, the Canadian summer bottom trawl survey data have been used exclusively to calibrate the SPA.

The survey data have been treated in various ways to account for the presence of sets with unusually large catches. In particular, these sets have been either removed or a Winsorizing technique employed to reduce their disproportionate impact on estimates of central tendency. O'Boyle and Wallace (1986) investigated the use of the delta distribution (Pennington, 1983) on this data set. The Statistics, Sampling and Survey Subcommittee (SSSS) of CAFSAC investigated the uses of this distribution and concluded that at low sample sizes ($n = 4$), the delta mean was very sensitive to the assumption that the population was indeed following the delta distribution. It stated that the arithmetic mean was not as sensitive to these assumptions and recommended its use unless the form of the distribution could be determined. Given the low sampling rates in the Canadian summer bottom trawl survey, the use of the arithmetic mean is preferred and was used in this assessment.

Tuning of the SPA has been effected using a variety of procedures. Initial efforts used geometric mean regressions between SPA abundance and the survey index, generally aggregated across age groups. More recently, age by age tuning has been conducted, using as criteria, maximization of the correlation coefficient (r) and attainment of a zero origin for the age specific calibration plots. Mohn (1983) showed that examination of the residuals of the last point would provide a more appropriate criterion. However, this places a lot of weight on the accuracy of the current year's data. To circumvent this problem, Doubleday (1981) had earlier used a combination of age-by-age relationships and examination of cohort-by-cohort fits to these relationships to establish current year fishing mortality, in a procedure called "Survivor". This procedure was used during 1981 - 86 to supplement the age-by-age tuned cohort analysis. In 1985, the principle of the examination of cohort-specific residuals was incorporated into the calibration of the cohort analysis. This procedure has remained unchanged since then although the underlying form of the calibration model (linear or curvilinear; additive or multiplicative errors) has been modified. Use of the survivor method was discontinued in this assessment.

The 1962 - 87 stock size was calculated and compared to estimates made by O'Boyle and Wallace (1986). Yield at $F_{0.1}$ and F_{50} (50% rule) was predicted for 1988.

Trends in Reported Landings

Annual Trends

The long-term (1930-1983) average annual catch for 4X haddock is about 20,000 t. This level has rarely been maintained for long periods. Prior to the mid-1960's, Canadian and US fishing vessels landed 15,000-20,000 t annually. In 1965-66, the Canadian offshore fleet expanded rapidly and displaced US fishing effort. As well, landings by the USSR substantially increased. This caused a rapid increase in reported landings, to levels never before attained by the resource (Figure 1). Much of this yield was supported by the 1963 year class, which has been shown (O'Boyle and Wallace, 1986) to be an order of magnitude larger than the average long term recruitment level.

Recruitment during the late 1960's was below average and could not support the high exploitation rates. ICNAF, the contemporary management body, was concerned with this situation and as a result imposed in 1970, the first quotas ever used by an international fisheries management organization. At the same time, the spawning grounds were closed to fishing during the spawning season.

Landings decreased in the early 1970's to levels above those seen prior to 1960. They remained low until 1975, just prior to extension of Canada's 200 mile limit in 1977. After this, landings increased rapidly, due to a recovery in the stocks as well as expansion of fleet size, fueled by federal and provincial vessel loan programs. A peak in catch was reached in 1981 and has declined to the 15,000 t level observed in 1985-86 (Table 1).

During the 1977-81 period, when fleet size was expanding and the TAC's were restrictive, misreporting and discarding became a significant problem. Reports of this occurring are based on discussions with fishermen and protection officers and have not been quantified. No estimate of misreporting and/or discarding are incorporated into this assessment.

The recent regulatory history of the stock is given in tables 2a and 2b. The quotas have been applied increasingly to smaller and smaller fleet units. In 1986-87, the allocation to mobile gear less than 65 ft. was also split by season. Table 2b illustrates that under the same allocation system, the 1987 fishery was slightly ahead of that of 1986.

Canadian Fishery by Gear Type, Tonnage Class and Unit Area

During the mid-1960's, activity by the mobile gear greater than 100 ft. represented the predominant amount of fishing effort directed on the resource. This fleet restricted its fishing effort to unit areas 4Xm-p (Figure 2), leaving the Bay of Fundy to the smaller draggers. The large vessel fleet substantially reduced its fishing operations in 4X in 1984 (Table 3), because of declining catch rates. As stated earlier, the mobile gear fleet less than 65 ft. increased in size during 1977-81 and became the dominant gear component by 1980 (Table 3).

This fleet operates throughout 4X but particularly in unit areas 4Xq-r during the latter part of the year. In 1986-87, this fleet increased its operations in 4VW, again due to declining catch rates in 4X.

The fixed gear, mostly longliner, fleet operates in unit areas 4Xm-p. Effort by this fleet has not changed dramatically. In recent years, there have been reports of large concentrations of dogfish which would tend to lower the catch rates of this fleet. This problem has decreased somewhat in 1986-87.

Age Composition of the Commercial Catch

Sampling Intensity

Since 1970, sampling for this resource has been generally good (Table 3). Prior to 1981, however, sampling was weighted heavily toward the offshore fleet and consequently sampling of inshore fleet catches was scanty. Since that time, sampling has been excellent with rates of one sample per 200-300 t landed attained. Sampling has been particularly good since 1985.

Construction of 1962-87 Catch-at-age

Construction of the catch-at-age matrix prior to 1978 is discussed by O'Boyle (1981). No adjustment has been made for the USSR catches in the mid-1960's. The length frequency of these landings was assumed to be the same as that for Canada. This is probably an inappropriate assumption and efforts are presently underway to rectify this. However, this should be kept in mind when analyzing the present data set for recruitment studies.

Construction of the 1978-85 catch at age is discussed by O'Boyle and Wallace (1986). The 1986 catch at age was constructed using the sampling stratum-sample availability table given in Table 4a. That for the first half of 1987 is given in Table 4b. This is in keeping with the practice followed since 1982.

For 1979-86, only Canadian sampling was used as no foreign sampling data were available. No correction for discarding and/or misreporting has been applied.

Trends in age composition

The catch and weight at age for 1962-86 are given in Table 5a. The pulse of the 1963 year-class through the fishery during the late 1960's is particularly evident. In recent years, no one year-class has supported the fishery. In 1986, the 1980-82 year-classes have been the dominant contributors. In 1987 (Table 5b), it is the 1981-83 year-classes that contribute the most to the landings. Consistently ages 4-6 have contributed the most numbers and biomass to the reported landings.

The 1986 observed catch-at-age contained substantially more younger fish than that projected by O'Boyle and Wallace (1986) (Table 6, Figure 3). The difference is particularly striking for the 1982 year-class which showed up stronger than expected.

The 1986 weights at age compare favourably to the long term (1962-86) average for the fishery (Figure 4). Certainly the recent declining trend in weights appears to have stopped.

An indication of the level of exploitation sustained by a resource can be obtained from examining trends in the average size and age of a fish in the catch. These parameters fluctuate dramatically due to recruitment pulses into the population. Nevertheless, the long term average level of these parameters is related to the exploitation rate. Trends in the 1962-86 estimates for average weight and age are given in Figure 5, along with the levels expected in populations exploited at $F_{0.1}$ and F_{max} . Since 1973, the average age in the catch has not reached above that expected for an F_{max} exploited stock. The average age has been stable around 4.75 since 1977. The average weight has also been generally below the F_{max} level and has exhibited a declining trend since 1982. These trends suggest high exploitation rates in the 4X haddock fishery.

Stock Abundance Trends

Commercial Catch Rates

Commercial catch rate indices were developed for various fleet sectors operating in 4X on a time and space basis (Table 7). Only trips where the main species was defined as being haddock were used. The catch rate was calculated as:

$$U_t = \frac{\sum Y_t}{\sum f_t} \quad (1)$$

where U_t is the catch rate in year t

Y_t is the catch biomass in year t for trips where effort has been reported and where haddock comprised the largest proportion of the catch for the trip (main species).

f_t is the effort in year t for trips where effort has been reported and where haddock comprised the largest proportion of the catch for the trip (main species).

A multiplicative model (Gavaris, 1980) incorporating all gears was not employed since the assumption of equivalent partial recruitment to all gears would be violated. Age structured multiplicative models (O'Boyle et al, 1984) are a possibility which remains to be explored.

Trends in the various catch rate series are provided in figures 6, 7 and 8. Those for the tonnage class 2-3 draggers all show a declining trend since 1979.

The catch rates for the tonnage class 4-5 vessels are relatively smooth in the first quarter but erratic in the second quarter. The latter trend is understandable given that the commercial fishing for this gear sector is exploited predominately in the first and fourth quarter of the year. The first quarter catch rates in 1986-87 dramatically increase to levels never before seen in the fishery. Examination of the raw data disclosed that information for only one trip (Cape John during 4 February - 5 March 1986) was present on the file. The trip had been particularly successful but fishing in 4X was discontinued for other fleet operation reasons. Although an investigation of the 1987 data was not undertaken it is likely that a similar event occurred. For this reason the 1986-87 catch rates cannot be compared to the historical time series.

The longliner catch rates (Figure 8) have been decreasing since 1984. Before then, the catch rates had been very erratic.

Three catch rate series for 1970-87 were chosen for further examination. All are reflective of catch rates at the time and location of most importance to their fleet specific fishery. The three are: 1) otter trawlers, TC 2-3, 1st quarter, 4Xm-p, 2) otter trawlers, TC 2-3, 3rd quarter, 4Xq-r and 3) otter trawlers, TC 4-5, 1st quarter, 4Xm-p. These catch rates were partitioned among age group using the port sampling data used to construct the catch at age matrix. The catch at age for these three fisheries is provided in Table 8, while the resulting catch rates, by age groups 2-5 (partially recruited) and 6-9 (fully recruited) are shown in Figure 9 and Table 9.

The spring fisheries show dramatic declines in the age 2-5 population since 1980. The trends in the older animals are not as clear.

Groundfish Bottom Trawl Surveys

Standardized Canadian summer bottom trawl surveys have been conducted in 4X since 1970. The vessel used during 1970-81 was the A. T. Cameron. In 1982 the Lady Hammond was used and in 1983 and 1984, the Alfred Needler was used. In the 1983 and 1984 assessments, discussion of conversion factors was extensive. In 1985, a conversion factor of $1.22 \times \text{ATC} = \text{LH} = \text{AN}$ was used. The SSS Subcommittee of CAFSAC met to decide the appropriate conversion factors and recommended that the A. T. Cameron values be adjusted upward by a factor of 1.2 ie. $1.2 \times \text{ATC} = \text{LH} = \text{AN}$. This was done here.

The arithmetic parameters for the summer survey data are given in Table 10.

Age 0+ abundance increased during 1977-80 and has declined gradually since. Age 2-5 abundance appears to have undergone a gradual increase since 1977 with a small decline in 1986 and a sharp decline in 1987. Age 6-9 abundance has gradually declined since 1977 (Figure 10).

Two recruitment trends were developed using the survey data set. These were calculated by first standardizing the age 1, 2 and 3 catch rates to the age specific 1970-87 average. Then the age 1 and 3 estimates on a cohort by cohort basis were averaged. The same was done for ages 2 and 3. The indices (Table 11, Figure 11) show a strong 1980 and 1982 year-class with relatively weak recruitment occurring in 1985 and 1986.

Sequential Population Analysis Methodology

The Calibration Data

The discrepancy between the survey and commercial catch rates noted in the last two assessments was once again observed. The former is providing a relatively more optimistic view of resource status compared to the latter. The survey series represents an unbiased but variable view of abundance. The commercial catch rate series represents a biased but relatively stable view of abundance. The latter series also suffers from the misreporting/discarding problems mentioned earlier. In addition, the fleets operating in 4X direct their operations in a multi-species fashion, making interpretation of catch rates even more problematical. For these reasons the previous practice of calibrating the SPA with survey data alone was continued.

The Calibration Model

As in O'Boyle and Wallace (1986) the following model was used as a basis of calibration:

$$U_{a,t} = (\alpha_a + q_a N_{a,t}) \cdot \epsilon_{a,t} \quad (2)$$

where $U_{a,t}$ is the survey catch numbers per tow at age a in year t

α_a is the age a intercept for the relationship.

q_a is the age a catchability of the survey or slope of the relationship.

$N_{a,t}$ is the mid-year age a , year t SPA abundance estimate.

$\epsilon_{a,t}$ is the error in the age a , year t estimate of the survey numbers per tow derived from the relationship.

The error term is assumed to be multiplicative rather than additive, based on analysis of the variance distribution in the data sets made by O'Boyle and Wallace (1986). Homogeneity of the variance is obtained through a log transform.

The survey catch rate was chosen as the dependent variable based on the observation that the variance in this data set is considerably higher than that in the catch at age data set (O'Boyle and Wallace, 1986).

The formulation requires use of a nonlinear least squares procedure. The algorithm used was a method of linearization which is based on the Newton-Raphson numerical technique (Draper and Smith, 1966).

The Calibration Methodology

The software used in calibration is given in Annex 1. The function AUTOSPANONLR is the main routine which calls INPUT for inputting all the basic matrices. ITERCOHORT (which calls PARTIAL), INTERFACERES and NONLR (which call TAYLOR and FCUM) represent the main calibration routines. The output are controlled by PLOTRESRVSPA (which calls PRINTRVSPA) and OUTPUT.

An initial run of SPA is conducted and equation 2 fit to the age 2 to 7 data, on an age by age basis, for the 1970-83 period. The 1984-87 data are not used for it is strongly influenced by the SPA input assumptions. The residuals for ages 2-7, 1970-87 are calculated using the age specific relationships. Starting at age 7, the 1987 fishing mortality which minimizes the sum of squares of the residuals along the 1980 cohort is chosen for the final 1987 exploitation rate. This fishing mortality is applied to ages 8 to 13+, thus implicitly assuming a flat-topped partial recruitment ogive. Once the age 7 estimate is obtained, the analysis is repeated for age 6 and the fishing mortality which minimizes the sum of squares of the 1981 year-class selected. This continues in a sequential manner for all ages down to age 2. This procedure is identical to that described by O'Boyle and Wallace (1986).

The age 12 and 13+ fishing mortality was calculated as

$$F = \frac{\sum_{a=1}^{13+} C_{a,t}}{\sum_{a=1}^{13+} \bar{N}_{a,t} \cdot PR_{a,t}} \quad (3)$$

where $PR_{a,t}$, the partial recruitment pattern in year t is calculated by dividing the mean F for ages 7 through 10 into that for age 1 to 13+ on a year by year basis.

As in the previous assessment, the 1987 first half of the year catch at age was estimated and included in the catch-at-age matrix to allow use of the 1987 survey data. No age 2 haddock were caught in the commercial fishery in the first half of 1987 (Table 5b). Consequently, the 1987 age 2 SPA could not be used to define the fishing mortality in that year on that age group. Nevertheless the age 2 relationship was used to define residuals for use in calibration of the older year-classes.

Results and Discussion

Model fit

The population and fishing mortality matrices for 1962-87 along with the calibration diagnostics are given in Tables 12 and 13 and Figure 12. The 1987 fishing mortalities are for only the first half of 1987, expressed in terms of a full year.

The fit of the model to the data is not as good as in O'Boyle and Wallace (1986). The correlation coefficients (r) are low, as indicated by the scatter of points around the regression lines (Figure 12a). Trends in the residuals with $\ln$ observed x or $\ln$ predicted y are not evident, indicating that the model choice is appropriate. However, the fit of the 1980-82 year classes is poor, with residuals being high at young ages (2-5) and low at older ages (6+). The model is unable to fit the high survey catch rates observed for these year classes at age 4. The predicted 1985-87 abundance for ages 5 and 6 is opposite in trend to that observed in the survey data. Nevertheless, this is the best fit of the model to the data, given the calibration technique.

Population Trends

Trends in age one beginning of year numbers (Figure 13a, Table 14) show that recruitment in recent years has dropped below the long term (1962-86) geometric mean of 28641×10^3 . Recent year-class strengths are ranked as:

Year-Class	Numbers at age one ($\times 10^{-6}$)	Ranking
1980	36.9	strong
1981	29.1	above average
1982	50.5	strong
1983	27.8	above average
1984	19.5	weak
1985	20.0	weak

The estimates for the 1980, 1981 and 1984 year-classes are substantially lower than those of O'Boyle and Wallace (1986). The 1982 year-class continues to be estimated at approximately 50×10^6 fish.

Mid-year population biomass (Figure 13b, Table 15) shows an increase to above 90,000 t during 1971-79 followed by a decline to 60,000 t by 1987. This year's evaluation of population biomass is substantially more pessimistic than that given by O'Boyle and Wallace (1986).

Finally, the fully recruited fishing mortality (Figure 13c, Table 15) shows exploitation well in excess of the $F_{0.1}$ level and for most years since 1977 in excess of F_{max} . It is evident that this resource is being subjected to growth overfishing.

Yield Projections to 1988

Yield was projected to 1988 assuming two different fishing mortality scenarios in 1987. The first assumed $F_{0.1}$ or 0.25 while the second used the 50 percent rule, which states that the exploitation rate should be half way between the current F (0.55) and the $F_{0.1}$ level (0.25). This provides an F of 0.40 in 1988.

The projection input conditions are given in Table 16. The beginning of 1987 population numbers for ages 3-13+ were taken directly from the SPA. The age 1 and 2 estimates of 20,000 and 16375×10^3 respectively were based upon examination of the age 1 + 2 survey recruitment index in relation to the SPA age one numbers. The 1985 and 1986 recruitment indices were the third and second lowest observed in the time series, behind the 1970 year-class. The next highest recruitment occurs at about $20,000 \times 10^3$ fish. Until more information is available on the strength of the 1985 and 1986 year-classes, it was felt that the figure of $20,000 \times 10^3$ at age 1 for these two year-classes would represent a conservative estimate of year-class strength. For the 1985 year-class, this provided an estimate of 16375×10^3 at age 2.

The weights at age used for projection were the 1984-86 average. The partial recruitment is also the annual average for this period. The annual vectors before averaging were calculated assuming full recruitment at ages 6 and 7 and adjusting all PR values above one to one. This and the weight at age vector are given in Table 16.

Natural mortality was assumed to be 0.2. The size of the 1987 year-class at age one was assumed to be the geometric mean of 1962-86 or 28641×10^3 .

The projection results are given in Table 17. Yield in 1988 at $F_{0.1}$ is 8204 t and 12,384 t at F_{50} . This level of harvesting is not significantly different from that presented by O'Boyle and Wallace (1986) and further emphasizes the poor state of the resource.

Concluding Remarks

The assessment presented here assumes that the survey data are tracking population abundance. The high fishing mortalities observed recently appear to be more in line with what is observed in the commercial catch rate series. If the 1988 survey indicates the same low stock size observed in the 1987 survey, then

reasons will have to be found to explain why the survey data indicated high stock size relative to the commercial data for the 1980-85 period.

Whatever the trend in abundance historically, it is evident that the resource is currently at a very low level, almost as low as that observed in the early 1970's. The resource is being growth overfished but it is difficult to establish if recruitment overfishing is occurring as well.

References

- Doubleday, W.G. 1981. A method of estimating the abundance of survivors of an exploited fish population using commercial fishing catch at age and research vessel abundance indices. Can. Spec. Publ. Fish. Aquat. Sci. 58: 164-178.
- Draper, N.R. and H. Smith. 1966. Applied regression analysis. John Wiley and Sons, N.Y. 407 p.
- Gavaris, S. 1980. Use of a multiplicative model to estimate catch rates and effort from commercial data. Can. J. Fish. Aquat. Sci. 37: 2272-2275.
- Mohn, R.F. 1983. Effects of error in catch and effort data in tuning cohort analysis, with a postscript on logistic growth models. Can. Spec. Publ. Fish. Aquat. Sci. 66: 141-150.
- O'Boyle, R. 1981. An assessment of the 4X haddock stock for the 1962-80 period. CAFSAC Res. Doc. 81/24.
- O'Boyle, R., J. McMillan, and G. White. 1984. The 4X haddock resource: A problem in supply and demand. CAFSAC Res. Doc. 84/100.
- O'Boyle, R. and D. Wallace. 1986. An Evaluation of the population Dynamics of 4X haddock during 1962-85 with yield projected to 1987. CAFSAC Res. Doc. 86/98.
- Pennington, M. 1983. Efficient estimators of abundance, for fish and plankton surveys. Biometrics. 39: 281-286.

Annex I

Software used in calibration of 1987 Division 4X haddock assessment.

Table 1. Reported nominal catch (t round) of haddock from NAFO Division 4X (excluding unit area 4Xs) by country. The numbers in brackets represent the number of commercial samples collected in that year.

YEAR	CANADA (MQ)	CANADA (Nfld)	USA	USSR	SPAIN	OTHER	TOTAL	TAC
1970	15560 (26)	-	1638	2	370	12	17582	18000
1971	16067 (29)	-	654	97	347	1	17166	18000
1972	12391 (36)	-	409	10	470	1	13281	9000
1973	12535 (30)	-	265	14	134	6	12954	9000
1974	12243 (25)	-	660	35	97	-	13035	-
1975	15985 (56)	-	2111	39	7	2	18144	15000
1976	16293 (45)	-	972	-	95	5	17365	15000
1977	19555 (79)	-	1648	2	-	12	21217	15000
1978	25299 (62)	114	1135	2	-	27	26577	21500
1979	24275 (49)	268	70	3	-	15	24631	26000
1980	28209 (56)	71	257	38	-	37	28612	28000
1981	30148 (82)	117	466	-	-	15	30746	27850
1982	23201 (92)	28	854	-	-	4	24087	32000
1983	24428 (119)	44	494	17	-	7	24990	32000
1984	19402 (97)	23	206	-	-	-	19631	32000
1985	14901 (86)	-	252	-	-	1	14927	15000
1986	14984 (78)	-	403	103	-	-	15034	15000
1987 ¹	8583 (56)	-	-	-	-	-	8583	15000

Long-term averages:

1930 - 60 = 16854 t
 1961 - 83 = 24217 t
 1930 - 83 = 20127 t

1. 1 January - 30 June 1986

2. NAFO SCS DOC 86/22.

3. NAFO SCS DOC 87/20.

Table 2a. Recent Canadian fishery allocations and the respective reported catch (t) of 4X haddock. Information from Atlantic Quota Reports.

Year	Report Date	Fleet	Allocation	Reported [†] Catch	%	CLOSURE DATES
1976		All Vessels	13300	15715	118	
1977		All Vessels	13400	20220	151	
1978		All Vessels	21500	25518	119	
1979		vessels < 125'	17500	17949	103	
		vessels > 125'	8500	6471	76	
		<u>Total</u>	26000	24420		
1980		vessels < 125'	22500	23585	105	
		vessels > 125'	5500	5095	93	
		<u>Total</u>	28000	28680		
1981	31/12	vessels < 125'	22350	25102	112	24/10 - 31/12
		vessels > 125'	5500	5380	98	02/02 - 31/12
		<u>Total</u>	27850	30482		
1982	31/12	FG. < 65'	8850	8168	92	
		MG. < 65'	15000	12909	86	
		FG. 65-100'	100	124	124	23/05 - 31/12
		MG. 65-100'	1000	567	57	
		MG. > 100'	7050	2829	40	
		<u>Total</u>	32000	24597		
1983	31/12	FG. < 65'	9050	9179	104	
		MG. < 65'	15000	12991	87	
		FG. 65-100'	100	108	108	12/04 - 31/12
		MG. 65-100'	800	177	18	
		MG. > 100'	7050	2438	35	
		<u>Total</u>	32000	24893		
1984	31/12	FG. < 65'	8850	6958	79	
		MG. < 65'	15000	12359	82	
		FG. 65-100'	100	3	3	
		MG. 65-100'	1000	44	4	
		MG. > 100'	7050	648	9	
		<u>Total</u>	32000	20012		
1985	31/12	FG. < 65'	4000	4496	112	16/11 - 31/12
		MG. < 65'	10000	10214	102	13/08 - 31/12
		FG. 65-100'	100	1	1	
		MG. 65-100'	100	61	61	
		MG. > 100'	800	541	68	
		<u>Total</u>	15000	15313		

Table 2a. (Continued)

Year	Report Date	Fleet	Allocation	Reported ¹ Catch	%	CLOSURE DATES
1986	31/12	FG. < 65'	5000	5446	109	
		MG. < 65'	1/1-30/4 2700			13/03
			1/5-31/8 4000			18/07
			1/9-31/12 2300	9202	102	
		FG. 65-100'	100	0	0	
		MG. 65-100'	100	118	118	15/02 , 15/11
		MG. > 100'	800	680	85	
		<u>Total</u>	15000	15446		
1987	26/09	FG. < 65'	5000	2944	59	
		MG. < 65'	1/1-30/4 2700			
			1/5-31/8 4000			
			1/9-31/12 2300	6299	70	
		FG. 65-100	100	49	49	
		MG. 65-100	100	118	118	
		MG. > 100	800	449	56	
		<u>Total</u>	15000	9859	66	

¹ These figures are based on haul information and thus are unofficial and not comparable to those in Table 1.

Table 2b. Trends in reported catch from Atlantic Quota Report to third week in April during 1982-87.

Fleet	1982	1983	1984	1985	1986	1987
FG < 65'	5146	4918	4397	2486	2631	2944
MG < 65'	8162	9057	8463	7457	5694	6299
FG 65-100'	119	171	2	1	0	49
MG 65-100'	451	185	33	46	110	118
MG > 100'	2179	2171	468	543	117	449

Table 3. Reported nominal catch (t round) of haddock from NAFO Division 4X (excluding unit areas 4Xs) landed in the Maritimes split by tonnage class and gear type. The numbers in brackets represent the mean weight landed per age/size sample collected.

1970-1987 sample corrected

Year	Tonnage Class				Total
	1 - 3		4+		
	MG	FG	MG	FG	
1970	4894 (979)	4049	6500 (295)	114	15559
71	4289 (715)	3973 (1006)	7712 (454)	93	16067
72	2741 (914)	4837 (1491)	4750 (238)	63	12391
73	1822 (304)	6415 (635)	4228 (302)	70	12535
74	3949 (790)	6615 (1120)	1623 (325)	56	12243
75	6091 (338)	5465 (1163)	4409 (457)	26	15991
76	4348 (1087)	5750 (1345)	6144 (486)	52	16294
77	6185 (1031)	4878 (490)	8345 (130)	152	19560
78	9213 (9213)	7904 (1083)	8093 (169)	92	25302
79	9870 (4935)	5727 (1076)	8634 (262)	56	24287
1980	13050 (1266)	8038 (932)	7045 (310)	82	28215
81	14769 (348)	8840 (540)	6475 (950)	70	30156
82	11614 (235)	8597 (470)	2972 (309)	32	23216
83	12315 (209)	9580 (438)	2538 (181)	15	24446
84	11744 (202)	7102 (1008)	609 (87)	0	19455
85	9727 (171)	4724 (168)	566 (113)	1	15018
86	9201 (192)	5575 (192)	209 (209)	1	14985
87 ¹ .	5650 (145)	2555 (213)	377 (75)	0	

1. 1 January - 30 June 1987.

Table 4a. Summary of commercial sampling for the haddock fishery in 1986. The tons landed is followed by sampling information in parentheses. The first number represents the number of fish measured and the second the number of otoliths read. The boxes represent the aggregation used in age-length key formation.

Quarter	<u>Mobile Gear</u>			
	TC 1-3	4Xm-p TC 4+	TC 1-3	4Xq-r TC 4+
1	2568 (4136-271)	147 (289-31)	157 (240-26)	0
2	830 (2025-144)	20	1317 (2501-146)	0
3	794 (761-72)	14	2284 (1192-143)	1
4	642 (225-31)	27	609 (494-27)	0

	<u>Fixed Gear</u>			
	TC 1-3	4Xm-p TC 4+	TC 1-3	4Xq-r TC 4+
1	1964 (2474-189)	-	5	-
2	329 (1002-105)	-	32	-
3	1719 (2463-163)	-	62	-
4	1451 (1152-130)	-	13	-

Table 4b. Summary of commercial sampling for the haddock fishery in 1987. The tons landed is followed by sampling information in parentheses. The first number represents the number of fish measured and the second the number of otoliths read. The boxes represent the aggregation used in age-length key formation.

QuarterMobile Gear

	TC 1-3	4Xm-p	TC 4+	TC 1-3	4Xq-r	TC 4+
1	3058 (4749-266)		219 (1121-77)	78		-
2	2061 (3174-84)		158	453 (1132-55)		-
3	-		-	-		-
4	-		-	-		-

Fixed Gear

	TC 1-3	4Xm-p	TC 4+	TC 1-3	4Xq-r	TC 4+
1	2210 (1797-151)		-	7		-
2	299 (711-89)		-	39		-
3	-		-	-		-
4	-		-	-		-

3.

35.2

	1970	1971	1972	1973	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	2031	2032	2033	2034	2035	2036	2037	2038	2039	2040	2041	2042	2043	2044	2045	2046	2047	2048	2049	2050	2051	2052	2053	2054	2055	2056	2057	2058	2059	2060	2061	2062	2063	2064	2065	2066	2067	2068	2069	2070	2071	2072	2073	2074	2075	2076	2077	2078	2079	2080	2081	2082	2083	2084	2085	2086	2087	2088	2089	2090	2091	2092	2093	2094	2095	2096	2097	2098	2099	2100	2101	2102	2103	2104	2105	2106	2107	2108	2109	2110	2111	2112	2113	2114	2115	2116	2117	2118	2119	2120	2121	2122	2123	2124	2125	2126	2127	2128	2129	2130	2131	2132	2133	2134	2135	2136	2137	2138	2139	2140	2141	2142	2143	2144	2145	2146	2147	2148	2149	2150	2151	2152	2153	2154	2155	2156	2157	2158	2159	2160	2161	2162	2163	2164	2165	2166	2167	2168	2169	2170	2171	2172	2173	2174	2175	2176	2177	2178	2179	2180	2181	2182	2183	2184	2185	2186	2187	2188	2189	2190	2191	2192	2193	2194	2195	2196	2197	2198	2199	2200	2201	2202	2203	2204	2205	2206	2207	2208	2209	2210	2211	2212	2213	2214	2215	2216	2217	2218	2219	2220	2221	2222	2223	2224	2225	2226	2227	2228	2229	2230	2231	2232	2233	2234	2235	2236	2237	2238	2239	2240	2241	2242	2243	2244	2245	2246	2247	2248	2249	2250	2251	2252	2253	2254	2255	2256	2257	2258	2259	2260	2261	2262	2263	2264	2265	2266	2267	2268	2269	2270	2271	2272	2273	2274	2275	2276	2277	2278	2279	2280	2281	2282	2283	2284	2285	2286	2287	2288	2289	2290	2291	2292	2293	2294	2295	2296	2297	2298	2299	2300	2301	2302	2303	2304	2305	2306	2307	2308	2309	2310	2311	2312	2313	2314	2315	2316	2317	2318	2319	2320	2321	2322	2323	2324	2325	2326	2327	2328	2329	2330	2331	2332	2333	2334	2335	2336	2337	2338	2339	2340	2341	2342	2343	2344	2345	2346	2347	2348	2349	2350	2351	2352	2353	2354	2355	2356	2357	2358	2359	2360	2361	2362	2363	2364	2365	2366	2367	2368	2369	2370	2371	2372	2373	2374	2375	2376	2377	2378	2379	2380	2381	2382	2383	2384	2385	2386	2387	2388	2389	2390	2391	2392	2393	2394	2395	2396	2397	2398	2399	2400	2401	2402	2403	2404	2405	2406	2407	2408	2409	2410	2411	2412	2413	2414	2415	2416	2417	2418	2419	2420	2421	2422	2423	2424	2425	2426	2427	2428	2429	2430	2431	2432	2433	2434	2435	2436	2437	2438	2439	2440	2441	2442	2443	2444	2445	2446	2447	2448	2449	2450	2451	2452	2453	2454	2455	2456	2457	2458	2459	2460	2461	2462	2463	2464	2465	2466	2467	2468	2469	2470	2471	2472	2473	2474	2475	2476	2477	2478	2479	2480	2481	2482	2483	2484	2485	2486	2487	2488	2489	2490	2491	2492	2493	2494	2495	2496	2497	2498	2499	2500	2501	2502	2503	2504	2505	2506	2507	2508	2509	2510	2511	2512	2513	2514	2515	2516	2517	2518	2519	2520	2521	2522	2523	2524	2525	2526	2527	2528	2529	2530	2531	2532	2533	2534	2535	2536	2537	2538	2539	2540	2541	2542	2543	2544	2545	2546	2547	2548	2549	2550	2551	2552	2553	2554	2555	2556	2557	2558	2559	2560	2561	2562	2563	2564	2565	2566	2567	2568	2569	2570	2571	2572	2573	2574	2575	2576	2577	2578	2579	2580	2581	2582	2583	2584	2585	2586	2587	2588	2589	2590	2591	2592	2593	2594	2595	2596	2597	2598	2599	2600	2601	2602	2603	2604	2605	2606	2607	2608	2609	2610	2611	2612	2613	2614	2615	2616	2617	2618	2619	2620	2621	2622	2623	2624	2625	2626	2627	2628	2629	2630	2631	2632	2633	2634	2635	2636	2637	2638	2639	2640	2641	2642	2643	2644	2645	2646	2647	2648	2649	2650	2651	2652	2653	2654	2655	2656	2657	2658	2659	2660	2661	2662	2663	2664	2665	2666	2667	2668	2669	2670	2671	2672	2673	2674	2675	2676	2677	2678	2679	2680	2681	2682	2683	2684	2685	2686	2687	2688	2689	2690	2691	2692	2693	2694	2695	2696	2697	2698	2699	2700	2701	2702	2703	2704	2705	2706	2707	2708	2709	2710	2711	2712	2713	2714	2715	2716	2717	2718	2719	2720	2721	2722	2723	2724	2725	2726	2727	2728	2729	2730	2731	2732	2733	2734	2735	2736	2737	2738	2739	2740	2741	2742	2743	2744	2745	2746	2747	2748	2749	2750	2751	2752	2753	2754	2755	2756	2757	2758	2759	2760	2761	2762	2763	2764	2765	2766	2767	2768	2769	2770	2771	2772	2773	2774	2775	2776	2777	2778	2779	2780	2781	2782	2783	2784	2785	2786	2787	2788	2789	2790	2791	2792	2793	2794	2795	2796	2797	2798	2799	2800	2801	2802	2803	2804	2805	2806	2807	2808	2809	2810	2811	2812	2813	2814	2815	2816	2817	2818	2819	2820	2821	2822	2823	2824	2825	2826	2827	2828	2829	2830	2831	2832	2833	2834	2835	2836	2837	2838	2839	2840	2841	2842	2843	2844	2845	2846	2847	2848	2849	2850	2851	2852	2853	2854	2855	2856	2857	2858	2859	2860	2861	2862	2863	2864	2865	2866	2867	2868	2869	2870	2871	2872	2873	2874	2875	2876	2877	2878	2879	2880	2881	2882	2883	2884	2885	2886	2887	2888	2889	2890	2891	2892	2893	2894	2895	2896	2897	2898	2899	2900	2901	2902	2903	2904	2905	2906	2907	2908	2909	2910	2911	2912	2913	2914	2915	2916	2917	2918	2919	2920	2921	2922	2923	2924	2925	2926	2927	2928	2929	2930	2931	2932	2933	2934	2935	2936	2937	2938	2939	2940	2941	2942	2943	2944	2945	2946	2947	2948	2949	2950	2951	2952	2953	2954	2955	2956	2957	2958	2959	2960	2961	2962	2963	2964	2965	2966	2967	2968	2969	2970	2971	2972	2973	2974	2975	2976	2977	2978	2979	2980	2981	2982	2983	2984	2985	2986	2987	2988	2989	2990	2991	2992	2993	2994	2995	2996	2997	2998	2999	3000
--	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------	------

Table 5b. Landings numbers, weight, percent numbers, and percent weight at age of NAFO Division 4X (excluding 4Xs) haddock for the first half of 1987.

Age	Numbers $\times 10^3$	s^2	C.V.	Weight (kg)	% by Number	% by Weight
1	0	0	0	-	0	0
2	0	0	0	-	0	0
3	420	2139.500	0.110	.626	6.32	3.07
4	1783	13921.800	0.066	.912	26.85	18.98
5	2762	18472.900	0.049	1.284	41.59	41.37
6	1419	10085.000	0.071	1.742	21.37	28.85
7	206	1010.100	0.154	2.453	3.10	5.90
8	45	130.800	0.250	2.960	.68	1.55
9	2	1.000	0.500	4.095	.03	0.10
10	2	-	-	4.410	.03	0.10
11	2	2.000	0.707	3.030	.03	0.07
12	0	-	-	-	0	0
13+	0	-	-	-	0	0

Table 6. Comparison of 1986 projected catch numbers with that observed in the 1986 fishery.

Age	Observed		Projected	
	No.	%	No.	%
1	0	0	1	0
2	290	2	172	2
3	1170	10	644	6
4	4378	38	2116	21
5	3923	34	4090	40
6	1476	13	2329	23
7	246	2	555	5
8	116	1	154	2
9	40	-	37	1
10	28	-	19	0
11	9	-	9	0
12	4	-	7	0
13+	2	-	3	0
No. (000's)	11682	100.00	10136	100.00
Catch (t)	15043		15000	

Table 7. Commercial catch rates (tons per unit effort) of 4X haddock for various fleet sectors. For otter trawlers the unit of effort is in hours fishing; for longliners, the unit of effort is in thousands of lines fished. QTR is the year quarter. Below the quarter is indicated the unit areas of capture (see Figure 2).

Year	OTB-2 (TC 2-3)			OTB-2 (TC 4-5)		LL (TC 2-3)		
	QTR1	QTR2	QTR3	QTR1	QTR2	QTR1	QTR2	QTR3
	MNOP	MNOP	OR	MNOP	MNOP	MNOP	MNOP	MNOP
1968	.279	.362	.665	.531	.373	3.161	7.029	-
69	-	.273	.576	.635	.563	14.997	-	-
1970	.064	.374	.155	.332	.650	10.261	-	-
71	.190	.369	.078	.282	.380	11.424	11.970	8.022
72	.099	.286	.094	.308	.390	16.789	4.822	7.413
73	-	.308	.564	.290	.231	15.824	4.076	23.208
74	.257	.142	.092	.335	.315	15.783	10.305	7.099
75	-	.298	.221	.424	.411	13.865	12.424	6.430
76	.362	.204	.205	.433	.352	15.204	9.197	6.647
77	.509	.349	.299	.479	.451	15.782	9.041	14.355
78	.428	.690	.247	.551	.646	17.214	14.147	7.877
79	.522	.461	.210	.895	.491	17.819	9.683	8.088
1980	.559	.329	.217	.827	.236	13.665	10.974	9.609
81	.462	.442	.289	.880	.889	15.011	8.184	8.743
82	.351	.231	.253	.642	.477	13.485	8.780	7.990
83	.287	.265	.180	.605	.380	13.985	10.883	8.951
84	.256	.259	.182	.413	.493	15.343	11.131	7.049
85	.283	.266	.125	.474	.389	13.345	9.883	5.882
86	.230	.142	.189	.923	-	11.083	6.365	7.260
87	.249	.213	-	1.490	-	12.685	5.848	-

Table 8. Landings at age of 4X haddock by gear sector in 1st and 3rd quarter.

[illegible][illegible][illegible]

Table 9. Catch rate (no. per hour fished) at age for Canadian otter trawlers fishing 4X haddock in unit areas MNOP (A and B) and OR (C) in the 1st and 3rd quarters during 1970-86.

		MATCH RATES (10 PER HOUR) AT AGE FOR 0.9, 100.0 ON FIRST STARTED																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																																									
		1970	1971	1972	1973	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987	1988	1989	1990	1991	1992	1993	1994	1995	1996	1997	1998	1999	2000	2001	2002	2003	2004	2005	2006	2007	2008	2009	2010	2011	2012	2013	2014	2015	2016	2017	2018	2019	2020	2021	2022	2023	2024	2025	2026	2027	2028	2029	2030	2031	2032	2033	2034	2035	2036	2037	2038	2039	2040	2041	2042	2043	2044	2045	2046	2047	2048	2049	2050	2051	2052	2053	2054	2055	2056	2057	2058	2059	2060	2061	2062	2063	2064	2065	2066	2067	2068	2069	2070	2071	2072	2073	2074	2075	2076	2077	2078	2079	2080	2081	2082	2083	2084	2085	2086	2087	2088	2089	2090	2091	2092	2093	2094	2095	2096	2097	2098	2099	2100	2101	2102	2103	2104	2105	2106	2107	2108	2109	2110	2111	2112	2113	2114	2115	2116	2117	2118	2119	2120	2121	2122	2123	2124	2125	2126	2127	2128	2129	2130	2131	2132	2133	2134	2135	2136	2137	2138	2139	2140	2141	2142	2143	2144	2145	2146	2147	2148	2149	2150	2151	2152	2153	2154	2155	2156	2157	2158	2159	2160	2161	2162	2163	2164	2165	2166	2167	2168	2169	2170	2171	2172	2173	2174	2175	2176	2177	2178	2179	2180	2181	2182	2183	2184	2185	2186	2187	2188	2189	2190	2191	2192	2193	2194	2195	2196	2197	2198	2199	2200	2201	2202	2203	2204	2205	2206	2207	2208	2209	2210	2211	2212	2213	2214	2215	2216	2217	2218	2219	2220	2221	2222	2223	2224	2225	2226	2227	2228	2229	2230	2231	2232	2233	2234	2235	2236	2237	2238	2239	2240	2241	2242	2243	2244	2245	2246	2247	2248	2249	2250	2251	2252	2253	2254	2255	2256	2257	2258	2259	2260	2261	2262	2263	2264	2265	2266	2267	2268	2269	2270	2271	2272	2273	2274	2275	2276	2277	2278	2279	2280	2281	2282	2283	2284	2285	2286	2287	2288	2289	2290	2291	2292	2293	2294	2295	2296	2297	2298	2299	2300	2301	2302	2303	2304	2305	2306	2307	2308	2309	2310	2311	2312	2313	2314	2315	2316	2317	2318	2319	2320	2321	2322	2323	2324	2325	2326	2327	2328	2329	2330	2331	2332	2333	2334	2335	2336	2337	2338	2339	2340	2341	2342	2343	2344	2345	2346	2347	2348	2349	2350	2351	2352	2353	2354	2355	2356	2357	2358	2359	2360	2361	2362	2363	2364	2365	2366	2367	2368	2369	2370	2371	2372	2373	2374	2375	2376	2377	2378	2379	2380	2381	2382	2383	2384	2385	2386	2387	2388	2389	2390	2391	2392	2393	2394	2395	2396	2397	2398	2399	2400	2401	2402	2403	2404	2405	2406	2407	2408	2409	2410	2411	2412	2413	2414	2415	2416	2417	2418	2419	2420	2421	2422	2423	2424	2425	2426	2427	2428	2429	2430	2431	2432	2433	2434	2435	2436	2437	2438	2439	2440	2441	2442	2443	2444	2445	2446	2447	2448	2449	2450	2451	2452	2453	2454	2455	2456	2457	2458	2459	2460	2461	2462	2463	2464	2465	2466	2467	2468	2469	2470	2471	2472	2473	2474	2475	2476	2477	2478	2479	2480	2481	2482	2483	2484	2485	2486	2487	2488	2489	2490	2491	2492	2493	2494	2495	2496	2497	2498	2499	2500	2501	2502	2503	2504	2505	2506	2507	2508	2509	2510	2511	2512	2513	2514	2515	2516	2517	2518	2519	2520	2521	2522	2523	2524	2525	2526	2527	2528	2529	2530	2531	2532	2533	2534	2535	2536	2537	2538	2539	2540	2541	2542	2543	2544	2545	2546	2547	2548	2549	2550	2551	2552	2553	2554	2555	2556	2557	2558	2559	2560	2561	2562	2563	2564	2565	2566	2567	2568	2569	2570	2571	2572	2573	2574	2575	2576	2577	2578	2579	2580	2581	2582	2583	2584	2585	2586	2587	2588	2589	2590	2591	2592	2593	2594	2595	2596	2597	2598	2599	2600	2601	2602	2603	2604	2605	2606	2607	2608	2609	2610	2611	2612	2613	2614	2615	2616	2617	2618	2619	2620	2621	2622	2623	2624	2625	2626	2627	2628	2629	2630	2631	2632	2633	2634	2635	2636	2637	2638	2639	2640	2641	2642	2643	2644	2645	2646	2647	2648	2649	2650	2651	2652	2653	2654	2655	2656	2657	2658	2659	2660	2661	2662	2663	2664	2665	2666	2667	2668	2669	2670	2671	2672	2673	2674	2675	2676	2677	2678	2679	2680	2681	2682	2683	2684	2685	2686	2687	2688	2689	2690	2691	2692	2693	2694	2695	2696	2697	2698	2699	2700	2701	2702	2703	2704	2705	2706	2707	2708	2709	2710	2711	2712	2713	2714	2715	2716	2717	2718	2719	2720	2721	2722	2723	2724	2725	2726	2727	2728	2729	2730	2731	2732	2733	2734	2735	2736	2737	2738	2739	2740	2741	2742	2743	2744	2745	2746	2747	2748	2749	2750	2751	2752	2753	2754	2755	2756	2757	2758	2759	2760	2761	2762	2763	2764	2765	2766	2767	2768	2769	2770	2771	2772	2773	2774	2775	2776	2777	2778	2779	2780	2781	2782	2783	2784	2785	2786	2787	2788	2789	2790	2791	2792	2793	2794	2795	2796	2797	2798	2799	2800	2801	2802	2803	2804	2805	2806	2807	2808	2809	2810	2811	2812	2813	2814	2815	2816	2817	2818	2819	2820	2821	2822	2823	2824	2825	2826	2827	2828	2829	2830	2831	2832	2833	2834	2835	2836	2837	2838	2839	2840	2841	2842	2843	2844	2845	2846	2847	2848	2849	2850	2851	2852	2853	2854	2855	2856	2857	2858	2859	2860	2861	2862	2863	2864	2865	2866	2867	2868	2869	2870	2871	2872	2873	2874	2875	2876	2877	2878	2879	2880	2881	2882	2883	2884	2885	2886	2887	2888	2889	2890	2891	2892	2893	2894	2895	2896	2897	2898	2899	2900	2901	2902	2903	2904	2905	2906	2907	2908	2909	2910	2911	2912	2913	2914	2915	2916	2917	2918	2919	2920	2921	2922	2923	2924	2925	2926	2927	2928	2929	2930	2931	2932	2933	2934	2935	2936	2937	2938	2939	2940	2941	2942	2943	2944	2945	2946	2947	2948	2949	2950	2951	2952	2953	2954	2955	2956	2957	2958	2959	2960	2961	2962	2963	2964	2965	2966	2967	2968	2969	2970	2971	2972	2973	2974	2975	2976	2977	2978	2979	2980	2981	2982	2983	2984	2985	2986	2987	2988	2989	2990	2991	2992	2993	2994	2995	2996	2997	2998	2999	3000	3001	3002	3003	3004	3005	3006	3007	3008	3009	3010	3011	3012	3013	3014	3015	3016	3017	3018	3019	3020	3021	3022	3023	3024	3025	3026	3027	3028	3029	3030	3031	3032	3033	3034	3035	3036	3037	3038	3039	3040	3041	3042	3043	3044	3045	3046	3047	3048	3049	3050	3051	3052	3053	3054	3055	3056	3057	3058	3059	3060	3061	3062	3063	3064	3065	3066	3067	3068	3069	3070	3071	3072	3073	3074	3075	3076	3077	3078	3079	3080	3081	3082	3083	3084	3085	3086	3087	3088	3089	3090	3091	3092	3093	3094	3095	3096	3097	3098	3099	3100	3101	3102	3103	3104	3105	3106	3107	3108	3109	3110	3111	3112	3113	3114	3115	3116	3117	3118	3119	3120	3121	3122	3123	3124	3125	3126	3127	3128	3129	3130	3131	3132	3133	3134	3135	3136	3137	3138	3139	3140	3141	3142	3143	3144	3145	3146	3147	3148	3149	3150	3151	3152	3153	3154	3155	3156	3157	3158	3159	3160	3161	3162	3163	3164	3165	3166	3167	3168	3169	3170	3171	3172	3173	3174	3175	3176	3177	3178	3179	3180	3181	3182	3183	3184	3185	3186	3187	3188	3189	3190	3191	3192	3193	3194	3195	3196	3197	3198	3199	3200	3201	3202	3203	3204	3205	3206	3207	3208	3209	3210	3211	3212	3213	3214	3215	3216	3217	3218	3219	3220	3221	3222	3223	3224	3225	3226	3227	3228	3229	3230	3231	3232	3233	3234	3235	3236	3237	3238	3239	3240	3241	3242	3243	3244	3245	3246	3247	3248	3249	3250	3251	3252	3253	3254	3255	3256	3257	3258	3259	3260	3261	3262	3263	3264	3265	3266	3267	3268	3269	3270	3271	3272	3273	3274	3275	3276	3277	3278	3279	3280	3281	3282	3283	3284	3285	3286	3287	3288	3289	3290	3291	3292	3293	3294	3295	3296	3297	3298	3299	3300	3301	3302	3303	3304	3305	3306	3307	3308	3309	3310	3311	3312	3313	3314	3315	3316	3317	3318	3319	3320	3321	3322	3323	3324	3325	3326	3327	3328	3329	3330	3331	3332	3333	3334	3335	3336	3337	3338	33

[illegible][illegible]

A.

3.



C.

11

Table 11. Recruitment indices for 4X haddock calculated from summer groundfish survey data set.

Year-Class	Age 1 + 2 Index	Age 2 + 3 Index
1968	-	0.394
1969	0.793	0.590
1970	0.017	0.028
1971	0.844	1.863
1972	1.195	1.145
1973	0.822	0.318
1974	0.653	1.913
1975	1.733	1.804
1976	0.575	0.633
1977	0.912	1.201
1978	0.379	0.573
1979	2.442	1.716
1980	3.548	1.783
1981	0.943	0.715
1982	1.346	2.012
1983	0.547	0.704
1984	0.743	0.512
1985	0.261	-
1986*	0.110	-

* Based on age one numbers per tow only.

Table 12. Cohort Analysis calibrated with survey catch per tow using Survey-SPA relationship assuming multiplicative error.

	1962	1963	1964	1965	1966	1967	1968	1969	1970	1971	1972	1973	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987
1	24986	91665	201009	141340	106774	17381	66633	14464	25294	6217	47285	44145	22719	40542	51656	20091	29147	27526	33956	26880	29147	56319	27011	27476	15823	15823
2	22402	20459	25550	145572	13115	6248	14220	6601	11842	20709	5090	36277	26006	17410	35720	31225	23016	27081	27514	27701	26194	23864	41367	22768	10431	10431
3	39711	24892	12104	61163	114677	10621	6713	11049	5195	6741	16242	4147	26802	28051	13710	31225	23016	27081	27514	27701	26194	23864	41367	22768	10431	10431
4	15226	44793	19195	12634	47444	63669	8020	5244	7222	3762	5693	10171	2293	17436	19483	9718	22931	24274	14651	19212	17608	14454	16875	14746	25431	14056
5	11366	14461	70173	12755	7441	2723	16266	21802	3166	1714	2442	1478	1407	2210	11240	12100	2973	17354	13045	9446	10102	8795	6879	9798	10191	16658
6	9587	7821	7046	16255	7441	2723	16266	21802	3166	1714	2442	1478	1407	2210	11240	12100	2973	17354	13045	9446	10102	8795	6879	9798	10191	16658
7	4087	5289	4744	2628	2043	2490	2597	1657	479	2016	1626	760	616	479	396	637	205	260	1400	914	702	630	575	226	140	203
8	1875	1694	1385	1127	1304	912	1912	1657	479	2016	1626	760	616	479	396	637	205	260	1400	914	702	630	575	226	140	203
9	1049	968	954	580	203	457	526	958	1172	272	728	1329	236	261	196	165	333	151	466	680	425	228	182	163	103	65
10	526	649	655	405	214	356	479	301	608	840	75	122	1402	455	60	43	55	176	103	255	160	110	43	15	10	6
11	202	253	253	157	80	146	300	201	273	91	340	524	130	59	714	224	14	14	3	7	49	33	41	26	2	6
12	316	185	92	265	300	327	368	201	125	344	250	17	16	34	341	147	115	31	25	19	26	41	17	2	3	0
13	167288	216024	363503	292453	223537	175515	119952	80504	70356	60496	87740	107620	103760	125061	140019	132758	136457	123439	171658	177772	117027	136609	118118	103643	163551	123276

9/ 7/87

FISHING MORTALITY

9/ 7/87

9/ 7/87

9/ 7/87

9/ 7/87

Table 13. Summary of diagnostics for final SPA calibration run

A. Regression Diagnostics

Age	a	$q_a \times 10^3$	R (1970-84)	R (1970-87)
2	-2.242	0.545	0.620	-
3	-2.071	0.687	0.680	0.725
4	0.003	0.469	0.581	0.568
5	-1.089	0.958	0.878	0.629
6	-0.593	1.125	0.740	0.588
7	0.145	0.798	0.896	0.879

B. In Residuals (obs-Pred)

AGE	YEAR																	
	1970	1971	1972	1973	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987
2	0.359	0.359	-0.007	-0.257	0.410	-0.586	-1.022	0.610	-0.641	-0.009	-0.253	0.949	0.830	-0.741	0.263	-0.277	0.461	-3.662
3	0.317	0.531	-0.730	-0.166	0.805	-1.070	-0.439	0.867	-0.526	0.045	0.140	-0.214	0.240	0.200	0.250	0.328	0.108	-0.464
4	-0.044	0.370	-0.373	-0.441	-0.311	0.134	-0.535	1.112	-0.949	0.214	0.365	0.291	-0.048	0.215	1.082	1.190	0.563	0.177
5	0.219	0.330	0.232	0.231	0.177	-0.574	0.122	0.398	-0.914	-0.416	0.471	0.409	0.274	-0.141	0.647	0.919	0.609	-0.847
6	0.120	0.422	-0.505	-0.398	0.419	-0.186	-0.209	0.656	-0.620	-0.037	0.154	-0.094	0.128	0.151	1.008	1.187	0.209	-0.670
7	-0.514	0.230	-0.335	-0.465	-0.157	-0.091	-0.531	0.264	-0.121	0.489	0.391	-0.037	0.803	0.073	0.984	0.865	0.082	-1.101

Table 14. Comparison of numbers (mill.) at age 1 generated by CAFSAC assessments since 1977.

Assessment	1970	1971	1972	1973	1974	1975	1976	1977	1978	1979	1980	1981	1982	1983	1984	1985	1986	1987
O'Boyle unpublished assessment #1 1977	30.8	7.0	60.4	49.3	28.9	33.4	-	-	-	-	-	-	-	-	-	-	-	-
O'Boyle unpublished assessment #2 1978	20.8	7.0	60.2	35.3	14.6	26.9	44.8	-	-	-	-	-	-	-	-	-	-	-
Res. Doc. 78/19	33.0	10.8	75.0	56.4	34.7	43.5	83.0	29.4	-	-	-	-	-	-	-	-	-	-
Res. Doc. 80/2	26.4	7.2	50.3	53.3	28.9	56.2	73.5	41.3	50.3	-	-	-	-	-	-	-	-	-
Res. Doc. 81/24	25.4	6.5	48.6	47.2	26.2	50.6	81.8	42.0	76.1	45.3	100.0	-	-	-	-	-	-	-
Res. Doc. 82/53	25.5	6.1	47.9	46.3	25.0	54.1	63.0	39.0	61.9	31.8	97.0	91.9	-	-	-	-	-	36
Res. Doc. 83/73	25.4	6.4	47.74	44.6	24.6	50.1	59.4	33.5	57.0	28.2	88.0	75.5	30.7	-	-	-	-	-
Res. Doc. 84/100	25.3	6.3	47.4	44.5	24.2	49.0	52.6	30.2	41.3	28.1	37.9	39.4	24.5	20.0	-	-	-	-
Res. Doc. 85/109	25.3	6.3	47.4	44.4	24.0	49.1	52.1	30.2	41.8	33.9	39.8	58.5	28.6	48.6	12.3	-	-	-
Res. Doc. 86/98	25.3	6.2	47.3	44.2	23.7	48.8	52.1	29.6	39.9	29.4	37.9	48.3	37.5	50.2	23.2	27.4	18.7	-
Present Document (Log)	25.3	6.2	47.3	44.1	23.7	48.6	51.6	29.1	39.1	27.5	34.1	36.9	29.1	50.5	27.8	19.5	20.0	20.0

Table 15. Comparison of population parameters generated by O'Boyle and Wallace (1986) and the current assessment.

Year	Mid-year Population Biomass (kt)		Fully Recruited ages (7-10) Fishing Mortality*	
	1986	1987	1986	1987
1970	59.6	59.6	0.394	0.394
71	49.9	49.9	0.633	0.633
72	53.2	53.2	0.516	0.516
73	58.0	58.0	0.473	0.473
74	60.1	60.0	0.592	0.593
75	71.3	71.2	0.433	0.433
76	79.7	79.4	0.697	0.698
77	81.9	81.4	1.021	1.024
78	93.3	92.4	0.603	0.608
79	94.6	92.9	0.480	0.486
80	88.3	85.6	0.567	0.578
81	87.1	80.5	0.702	0.727
82	84.7	73.4	0.713	0.768
83	80.2	67.4	0.932	1.088
84	78.3	64.1	0.916	1.498
85	81.0	64.7	0.422	0.991
86	80.6	62.4	0.387	0.513
87	-	60.3	-	0.549

* ages 6-7 for 1986 only

Table 16. 1987 population conditions used for projection purposes

Age	Beg. of year Numbers at age (000's)	Weight at age (kg) (1984-86)	Partial Recruitment (1984-86)
1	20000	0.250	0.0001
2	16375	0.534	0.020
3	12807	0.766	0.081
4	14056	1.054	0.270
5	16859	1.400	0.719
6	4794	1.914	1.000
7	1752	2.409	1.000
8	383	2.733	1.000
9	17	3.265	1.000
10	17	3.405	1.000
11	17	3.680	1.000
12	0	3.841	1.000
13	0	4.703	1.000

$M = 0.2$

$F_{0.1} = 0.25$

Age one recruitment (1962-86 geometric mean) = 28641×10^3

Table 17. Catch projection results using input data given in Table 16.

Year	Mid-year 1+ Population Biomass (t)	1+ Catch Biomass (t)	Fully Recruited Fishing Mortality
1987	61708	15000	0.549
1988 ($F_{0.1}$)	65448	8204	0.25
1988 (F_{50})	63246	12384	0.40




Figure 1. Long-term trends in 4X haddock landings, along with annual TAC since 1970.




Figure 2. Canadian fisheries statistical unit areas in NAFO Division 4X




Figure 3. Catch numbers (000's) at age for 4X haddock. Comparison of observed 1986 catch at age (—) with that projected, using 15,000 t in 1986, by O'Boyle and Wallace, 1986 (◊).




Figure 4. Weight at age (kg) of haddock in commercial catch samples in NAFO Division 4X, 1962-1986. Weights at age for 1986 (◆) and mean weights at age for 1962-86 (◻)





Figure 5. Age-size characteristics of landings of 4X haddock.

A. Average age of 4X haddock in landings.

B. Average weight (kg) of 4X haddock in landings.

Top and bottom line in each figure indicates levels of these parameters in populations harvested at $F_{0.1}$ and F_{max} respectively.





Figure 6. Trends in catch rates (T per hour fished) of haddock by other trawlers, 1965-1985, fishing in 1st quarter, 48 mph (A), 2nd quarter, 50 mph (B), and third quarter, 48 mph (C).

A.



B.




Figure 7. Trends in catch rates (t per hour fished) of haddock by otter trawlers, TC 4-5, fishing in 4X MNOP (A) during 1st and 2nd quarter.






Figure 8. Trends in catch rates (t per thousands of lines fished) of haddock by longliners, IC 2-3, fishing in 4X MBOP during 1st (A), 2nd (B) and 3rd (C) quarters.




Figure 9. Trends in catch rates (no. per hour fished) of haddock for ages 2-5 (A,C,E) and age 6-9 (B,D,F) by otter trawlers.

A-B: TC 2-3, 1st quarter, 4X MNOP.

C-D: TC 4-5, 1st quarter, 4X MNOP.

E-F: TC 2-3, 3rd quarter, 4X QR.






Figure 10. Survey arithmetic mean catch rates (#/tow) of haddock from 4X during 1970-87 for ages 2-5 (A), 6-9 (B) and all age groups combined (C). The solid line in each figure indicates the trend in the median smoothed estimates






Figure 11. Trend in age 1 + 2 and age 2 + 3 recruitment indices derived from Canadian summer survey data set.




Figure 12a. Relationships by age between survey numbers per tow and SPA mid-year numbers from cohort analysis calibrated with equation 2 in text. Solid line is predicted relationship. Number beside point indicates year-class.




Figure 12b. Relationships by age between LN survey numbers per tow and LN SPA mid-year numbers from cohort analysis calibrated with equation 2 in text. Solid line is predicted relationship.




Figure 12c. Trend in age-specific predicted (solid line) and observed survey numbers per tow derived from cohort analysis calibrated using equation 2 in text. Number beside point indicates year-class.




Figure 12d. Trend in LN residuals with time for age-by-age relationships of cohort analysis calibrated using equation 2 in text.




Figure 12e. Relationships by age between LN residuals and LN predicted survey numbers per tow for calibration equation 2 given in text.




Figure 12f. Relationship by age between LN residuals and LN observed survey numbers per tow for calibration equation 2 given in text.




Figure 13. Trends in age one population numbers (A), mid-year population biomass (B), and fully recruited fishing mortality (C) of the 4X haddock resource. All figures show the results of the current assessment (1987) in comparison to that of last year (1986).

